

Author Index

Adamec, E., Beermann, M.L. and Nixon, R.A. Calpain I activation in rat hippocampal neurons in culture is NMDA receptor selective and not essential for excitotoxic cell death (54) 35

Afify, E.A., Law, P.Y., Riedl, M., Elde, R. and Loh, H.H.

Role of carboxyl terminus of μ - and δ opioid receptor in agonist -induced downregulation (54) 24

Bader, M., see Baltatu, O. (54) 237 Baltatu, O., Lippoldt, A., Hansson, A., Ganten, D. and Bader, M. Local renin-angiotensin system in the pineal

gland (54) 237 Beach, T.G., see Cicchetti, F. (54) 1 Beermann, M.L., see Adamec, E. (54) 35

Besson, M.-J., see Mathieu-Kia, A.-M. (54) 141 Bowery, N.G., see Meoni, P. (54) 13 Bunnemann, B.H., see Meoni, P. (54) 13

Cagampang, F.R.A., Sheward, W.J., Harmar, A.J., Piggins, H.D. and Coen, C.W. Circadian changes in the expression of vasoactive intestinal peptide 2 receptor mRNA in the rat suprachiasmatic nuclei (54) 108 Cavallaro, S., see D'Agata, V. (54) 161

Cavalli, A., see Loh, H.H. (54) 321

Chen, A.C.-H., McDonald, B., Moss, S.J. and Gurling, H.M.D.

Gene expression studies of mRNAs encoding the NMDA receptor subunits NMDAR1, NMDAR2A, NMDAR2B, NMDAR2C, and NMDAR2D following long-term treatment with cis- and trans-flupenthixol as a model for understanding the mode of action of schizophrenia drug treatment (54) 92

Chen, Y.-F., see Loh, H.H. (54) 321 Chi, S., see McMahon, A. (54) 56 Cho, H., see Cho, S. (54) 74

Cho, S., Cho, H., Geum, D. and Kim, K. Retinoic acid regulates gonadotropin-releasing hormone (GnRH) release and gene expression in the rat hypothalamic fragments and GT1-1 neuronal cells in vitro (54) 74

Cicchetti, F., Lacroix, S., Beach, T.G. and Parent, A.

Calretinin gene expression in the human thalamus (54) 1

Copeland, N.G., see Kamata, T. (54) 219 Copeland, T.D., see Kamata, T. (54) 219

D'Agata, V. and Cavallaro, S. Functional and molecular expression of PACAP/VIP receptors in the rat retina (54)

David, S., Shoemaker, M. and Haley, B.E. Abnormal properties of creatine kinase in Alzheimer's disease brain: Correlation of reduced enzyme activity and active site photolabeling with aberrant cytosol-membrane partitioning (54) 276

Donohue, P.J., Shapira, H., Mantey, S.A., Hampton, L.L., Jensen, R.T. and Battey,

A human gene encodes a putative G protein-coupled receptor highly expressed in the central nervous system (54) 152

Elde, R., see Afify, E.A. (54) 24

Fan, F.S., Shen, H.-H., Tseng, W.-P., Chen, P.-M. and Tsai, S.-F. Molecular cloning and characterization of a human brain-specific gene implicated in neuronal differentiation (54) 113

Friberg, I.K., Young, A.B. and Standaert, D.G. Differential localization of the mRNAs for the pertussis toxin insensitive G-protein alpha sub-units Ga, G11, and G2 in the rat brain, and regulation of their expression after striatal deafferentation (54) 298

Furuya, S., see Futami, T. (54) 183

Futami, T., Hatanaka, Y., Matsushita, K. and Furuva, S. Expression of substance P receptor in the substantia nigra (54) 183

Ganten, D., see Baltatu, O. (54) 237 Garbay, B., see Kursula, P. (54) 252 German, D.C., see McMahon, A. (54) 56 Geum, D., see Cho, S. (54) 74 Gintzler, A.R., see Rivera, M. (54) 165 Gurling, H.M.D., see Chen, A.C.-H. (54) 92 Gustafsson, J.-Å., see Österlund, M. (54) 175

Haley, B.E., see David, S. (54) 276

Hamada, S., Senzaki, K., Hamaguchi-Hamada, K., Tabuchi, K., Yamamoto, H., Yamamoto, T., Yoshikawa, S., Okano, H. and Okado, N. Localization of 5-HT2A Receptor in rat cerebral cortex and olfactory system revealed by immunohistochemistry using two antibodies raised in rabbit and chicken (54) Hampton, L.L., see Donohue, P.J. (54) 152 Hansson, A., see Baltatu, O. (54) 237 Hara, Y., see Kamata, T. (54) 219 Harmar, A.J., see Cagampang, F.R.A. (54) 108 Hatanaka, Y., see Futami, T. (54) 183 Heape, A.M., see Kursula, P. (54) 252 Hibino, Y., see Katsura, M. (54) 124 Hibino, Y., see Ohkuma, S. (54) 133 Hirao, T., Morimoto, K., Yamamoto, Y., Watanabe, T., Sato, H., Sato, K., Sato, S., Yamada, N., Tanaka, K. and Suwaki, H. Time-dependent and regional expression of GABA transporter mRNAs following amygdala-kindled seizures in rats (54) 49 Huh, S.-O., see Won, J.-S. (54) 288

Iacopino, A.M., see McMahon, A. (54) 56 Imaizumi, K., see Kanazawa, K.-y. (54) 316 Imaizumi, K., see Tanimukai, H. (54) 212 Ishikawa, E., see Katsura, M. (54) 124

Hurd, Y.L., see Österlund, M. (54) 175

Jenkins, N.A., see Kamata, T. (54) 219 Jensen, R.T., see Donohue, P.J. (54) 152

Kamata, T., Subleski, M., Hara, Y., Yuhki, N., Kung, H.-f., Copeland, N.G., Jenkins, N.A., Yoshimura, T., Modi, W. and Copeland,

Isolation and characterization of a bovine neural specific protein (CRMP-2) cDNA homologous to unc-33, a C. elegans gene implicated in axonal outgrowth and guidance (54) 219

Kanazawa, K .- y ., Imaizumi, K ., Mori, T ., Honma, Y., Tojo, M., Tanno, Y., Yokoya, S., Niwa, S.-i., Tohyama, M., Takagi, T. and Wanaka, A.

Expression pattern of a novel death-promoting gene, DP5, in the developing murine nervous system (54) 316

Katsura, M., Ohkuma, S., Tsujimura, A., Xu, J., Hibino, Y., Ishikawa, E. and Kuriyama, K. Functional involvement of benzodiazepine receptors in ethanol-induced increases of diazepam binding inhibitor (DBI) and its mRNA in the mouse brain (54) 124

Kim, K., see Cho, S. (54) 74 Kim, Y.-H., see Won, J.-S. (54) 288 Kitani, T., see Sakagami, H. (54) 311 Kondo, H., see Sakagami, H. (54) 311



Author Index

Adamec, E., Beermann, M.L. and Nixon, R.A. Calpain I activation in rat hippocampal neurons in culture is NMDA receptor selective and not essential for excitotoxic cell death (54) 35

Afify, E.A., Law, P.Y., Riedl, M., Elde, R. and Loh, H.H.

Role of carboxyl terminus of μ - and δ opioid receptor in agonist -induced downregulation (54) 24

Bader, M., see Baltatu, O. (54) 237 Baltatu, O., Lippoldt, A., Hansson, A., Ganten, D. and Bader, M. Local renin-angiotensin system in the pineal

gland (54) 237 Beach, T.G., see Cicchetti, F. (54) 1 Beermann, M.L., see Adamec, E. (54) 35

Besson, M.-J., see Mathieu-Kia, A.-M. (54) 141 Bowery, N.G., see Meoni, P. (54) 13 Bunnemann, B.H., see Meoni, P. (54) 13

Cagampang, F.R.A., Sheward, W.J., Harmar, A.J., Piggins, H.D. and Coen, C.W. Circadian changes in the expression of vasoactive intestinal peptide 2 receptor mRNA in the rat suprachiasmatic nuclei (54) 108 Cavallaro, S., see D'Agata, V. (54) 161

Cavalli, A., see Loh, H.H. (54) 321

Chen, A.C.-H., McDonald, B., Moss, S.J. and Gurling, H.M.D.

Gene expression studies of mRNAs encoding the NMDA receptor subunits NMDAR1, NMDAR2A, NMDAR2B, NMDAR2C, and NMDAR2D following long-term treatment with cis- and trans-flupenthixol as a model for understanding the mode of action of schizophrenia drug treatment (54) 92

Chen, Y.-F., see Loh, H.H. (54) 321 Chi, S., see McMahon, A. (54) 56 Cho, H., see Cho, S. (54) 74

Cho, S., Cho, H., Geum, D. and Kim, K. Retinoic acid regulates gonadotropin-releasing hormone (GnRH) release and gene expression in the rat hypothalamic fragments and GT1-1 neuronal cells in vitro (54) 74

Cicchetti, F., Lacroix, S., Beach, T.G. and Parent, A.

Calretinin gene expression in the human thalamus (54) 1

Copeland, N.G., see Kamata, T. (54) 219 Copeland, T.D., see Kamata, T. (54) 219

D'Agata, V. and Cavallaro, S. Functional and molecular expression of PACAP/VIP receptors in the rat retina (54)

David, S., Shoemaker, M. and Haley, B.E. Abnormal properties of creatine kinase in Alzheimer's disease brain: Correlation of reduced enzyme activity and active site photolabeling with aberrant cytosol-membrane partitioning (54) 276

Donohue, P.J., Shapira, H., Mantey, S.A., Hampton, L.L., Jensen, R.T. and Battey,

A human gene encodes a putative G protein-coupled receptor highly expressed in the central nervous system (54) 152

Elde, R., see Afify, E.A. (54) 24

Fan, F.S., Shen, H.-H., Tseng, W.-P., Chen, P.-M. and Tsai, S.-F. Molecular cloning and characterization of a human brain-specific gene implicated in neuronal differentiation (54) 113

Friberg, I.K., Young, A.B. and Standaert, D.G. Differential localization of the mRNAs for the pertussis toxin insensitive G-protein alpha sub-units Ga, G11, and G2 in the rat brain, and regulation of their expression after striatal deafferentation (54) 298

Furuya, S., see Futami, T. (54) 183

Futami, T., Hatanaka, Y., Matsushita, K. and Furuva, S. Expression of substance P receptor in the substantia nigra (54) 183

Ganten, D., see Baltatu, O. (54) 237 Garbay, B., see Kursula, P. (54) 252 German, D.C., see McMahon, A. (54) 56 Geum, D., see Cho, S. (54) 74 Gintzler, A.R., see Rivera, M. (54) 165 Gurling, H.M.D., see Chen, A.C.-H. (54) 92 Gustafsson, J.-Å., see Österlund, M. (54) 175

Haley, B.E., see David, S. (54) 276

Hamada, S., Senzaki, K., Hamaguchi-Hamada, K., Tabuchi, K., Yamamoto, H., Yamamoto, T., Yoshikawa, S., Okano, H. and Okado, N. Localization of 5-HT2A Receptor in rat cerebral cortex and olfactory system revealed by immunohistochemistry using two antibodies raised in rabbit and chicken (54) Hampton, L.L., see Donohue, P.J. (54) 152 Hansson, A., see Baltatu, O. (54) 237 Hara, Y., see Kamata, T. (54) 219 Harmar, A.J., see Cagampang, F.R.A. (54) 108 Hatanaka, Y., see Futami, T. (54) 183 Heape, A.M., see Kursula, P. (54) 252 Hibino, Y., see Katsura, M. (54) 124 Hibino, Y., see Ohkuma, S. (54) 133 Hirao, T., Morimoto, K., Yamamoto, Y., Watanabe, T., Sato, H., Sato, K., Sato, S., Yamada, N., Tanaka, K. and Suwaki, H. Time-dependent and regional expression of GABA transporter mRNAs following amygdala-kindled seizures in rats (54) 49 Huh, S.-O., see Won, J.-S. (54) 288

Iacopino, A.M., see McMahon, A. (54) 56 Imaizumi, K., see Kanazawa, K.-y. (54) 316 Imaizumi, K., see Tanimukai, H. (54) 212 Ishikawa, E., see Katsura, M. (54) 124

Hurd, Y.L., see Österlund, M. (54) 175

Jenkins, N.A., see Kamata, T. (54) 219 Jensen, R.T., see Donohue, P.J. (54) 152

Kamata, T., Subleski, M., Hara, Y., Yuhki, N., Kung, H.-f., Copeland, N.G., Jenkins, N.A., Yoshimura, T., Modi, W. and Copeland,

Isolation and characterization of a bovine neural specific protein (CRMP-2) cDNA homologous to unc-33, a C. elegans gene implicated in axonal outgrowth and guidance (54) 219

Kanazawa, K .- y ., Imaizumi, K ., Mori, T ., Honma, Y., Tojo, M., Tanno, Y., Yokoya, S., Niwa, S.-i., Tohyama, M., Takagi, T. and Wanaka, A.

Expression pattern of a novel death-promoting gene, DP5, in the developing murine nervous system (54) 316

Katsura, M., Ohkuma, S., Tsujimura, A., Xu, J., Hibino, Y., Ishikawa, E. and Kuriyama, K. Functional involvement of benzodiazepine receptors in ethanol-induced increases of diazepam binding inhibitor (DBI) and its mRNA in the mouse brain (54) 124

Kim, K., see Cho, S. (54) 74 Kim, Y.-H., see Won, J.-S. (54) 288 Kitani, T., see Sakagami, H. (54) 311 Kondo, H., see Sakagami, H. (54) 311 Kovács, K.J., Arias, C. and Sawchenko, P.E.
Protein synthesis blockade differentially affects the stress- induced transcriptional activation of neuropeptide genes in parvocellular neurosecretory neurons (54) 85
Kuiper, G.G.J.M., see Österlund, M. (54) 175
Kung, H.-f., see Kamata, T. (54) 219
Kuriyama, K., see Katsura, M. (54) 124
Kuriyama, K., see Ohkuma, S. (54) 133
Kursula, P., Lehto, V.-P., Garbay, B., Cassagne, C. and Heape, A.M.
Expression of the amino acid dimorphism in the small myelin-associated glycoprotein cytoplasmic domain in rat peripheral nerves during postnatal development (54) 252

Lacroix, S., see Cicchetti, F. (54) 1
Law, P.Y., see Afify, E.A. (54) 24
Lehto, V.-P., see Kursula, P. (54) 252
Lippoldt, A., see Baltatu, O. (54) 237
Liu, H.-C., see Loh, H.H. (54) 321
Loh, H.H., Liu, H.-C., Cavalli, A., Yang, W.,
Chen, Y.-F. and Wei, L.-N.

μ Opioid receptor knockout in mice: effects
on ligand-induced analgesia and morphine
lethality (54) 321

Mantey, S.A., see Donohue, P.J. (54) 152
Mathieu-Kia, A.-M. and Besson, M.-J.
Repeated administration of cocaine, nicotine
and ethanol: effects on preprodynorphin,
preprotachykinin A and preproenkephalin
mRNA expression in the dorsal and the
ventral striatum of the rat (54) 141
McDonald, B., see Chen, A.C.-H. (54) 92
McGinty, J.F., see Wang, J.Q. (54) 262
McMahon, A., Wong, B.S., Iacopino, A.M.,
Ng, M.C., Chi, S. and German, D.C.
Calbindin-D_{28k} buffers intracellular calcium
and promotes resistance to degeneration in
PC12 cells (54) 56
Meoni, P., Mugnaini, M., Bunnemann, B.H.,

Meoni, P., Mugnaini, M., Bunnemann, B.H., Trist, D.G. and Bowery, N.G. [³H]MK-801 binding and the mRNA for the NMDAR1 subunit of the NMDA receptor are differentially distributed in human and rat forebrain (54) 13

Mhatre, M.C. and Ticku, M.K.
Caloric restriction retards the aging associated changes in γ-aminobutyric acid_A receptor gene expression in rat cerebellum (54) 270

Mijnster, M.J., Galis-de Graaf, Y. and Voorn, P.
Serotonergic regulation of neuropeptide and glutamic acid decarboxylase mRNA levels in the rat striatum and globus pallidus: studies with fluoxetine and DOI (54) 64

Modi, W., see Kamata, T. (54) 219 Mori, T., see Kanazawa, K.-y. (54) 316 Morimoto, K., see Hirao, T. (54) 49 Morley, B.J. ARIA is heavily expressed in rat peripheral

ARIA is heavily expressed in rat peripheral auditory and vestibular ganglia (54) 170 Mugnaini, M., see Meoni, P. (54) 13

Ng, M.C., see McMahon, A. (54) 56 Niwa, S.-i., see Kanazawa, K.-y. (54) 316 Nixon, R.A., see Adamec, E. (54) 35

Ohkuma, S., Katsura, M., Hibino, Y., Xu, J., Shirotani, K. and Kuriyama, K. Multiple actions of nitric oxide on voltage-dependent Ca²⁺ channels in mouse cerebral cortical neurons (54) 133
Okado, N., see Hamada, S. (54) 199
Okano, H., see Hamada, S. (54) 199
Okuno, S., see Sakagami, H. (54) 311
Österlund, M., Kuiper, G.G.J.M., Gustafsson, J.-Å. and Hurd, Y.L.
Differential distribution and regulation of estrogen receptor-α and -β mRNA within the female rat brain (54) 175

Piggins, H.D., see Cagampang, F.R.A. (54) 108 Przewłocka, B., see Ziółkowska, B. (54) 243 Przewłocki, R., see Ziółkowska, B. (54) 243

Rashid, A.J. and Dunn, R.J.
Sequence diversity of voltage-gated potassium channels in an electric fish (54) 101
Rivera, M. and Gintzler, A.R.
Differential effect of chronic morphine on

Differential effect of chronic morphine on mRNA encoding adenylyl cyclase isoforms: relevance to physiological sequela of tolerance/dependence (54) 165

Sakagami, H., Saito, S., Kitani, T., Okuno, S., Fujisawa, H. and Kondo, H.
 Localization of the mRNAs for two isoforms of Ca²⁺/calmodulin-dependent protein kinase kinases in the adult rat brain (54) 311
 Sato, K., see Hirao, T. (54) 49

Sato, K., see Hirao, T. (54) 49
Sato, S., see Hirao, T. (54) 49
Sawchenko, P.E., see Kovács, K.J. (54) 85
Senzaki, K., see Hamada, S. (54) 199
Shapira, H., see Donohue, P.J. (54) 152
Shen, H.-H., see Fan, F.S. (54) 113
Sheward, W.J., see Cagampang, F.R.A. (54) 108

Shirotani, K., see Ohkuma, S. (54) 133 Shoemaker, M., see David, S. (54) 276 Song, D.-K., see Won, J.-S. (54) 288 Standaert, D.G., see Friberg, I.K. (54) 298 Subleski, M., see Kamata, T. (54) 219 Suh, H.-W., see Won, J.-S. (54) 288 Suwaki, H., see Hirao, T. (54) 49 Szél, A., see Von Schantz, M. (54) 327 Tanaka, K., see Hirao, T. (54) 49
Tanimukai, H., Imaizumi, K., Kudo, T.,
Katayama, T., Tsuda, M., Takagi, T., Tohyama, M. and Takeda, M.
Alzheimer-associated presenilin-1 gene is
induced in gerbil hippocampus after transient ischemia (54) 212
Ticku, M.K., see Mhatre, M.C. (54) 270
Tohyama, M., see Kanazawa, K.-y. (54) 316
Tohyama, M., see Tanimukai, H. (54) 212
Tojo, M., see Kanazawa, K.-y. (54) 316
Trist, D.G., see Meoni, P. (54) 13
Tsai, S.-F., see Fan, F.S. (54) 113
Tseng, W.-P., see Fan, F.S. (54) 113
Tsuda, M., see Tanimukai, H. (54) 212

Tabuchi, K., see Hamada, S. (54) 199

Takagi, T., see Kanazawa, K.-y. (54) 316

Takagi, T., see Tanimukai, H. (54) 212

Takeda, M., see Tanimukai, H. (54) 212

Van Veen, T., see Von Schantz, M. (54) 327 Von Schantz, M., Szél, A., Van Veen, T. and Farber, D.B. Cloning of a cyclic GMP phosphodiesterase y subunit from the ground squirrel retina (54) 327

Tsujimura, A., see Katsura, M. (54) 124

Wanaka, A., see Kanazawa, K.-y. (54) 316
Wang, J.Q. and McGinty, J.F.
Metabotropic glutamate receptor agonist increases neuropeptide mRNA expression in rat striatum (54) 262

Wei, L.-N., see Loh, H.H. (54) 321
Won, J.-S., Song, D.-K., Kim, Y.-H., Huh, S.-O. and Suh, H.-W.
The stimulation of rat astrocytes with phorbol-12-myristate-13-acetate increases the proenkephalin mRNA: involvement of proto-oncogenes (54) 288

Xu, J., see Katsura, M. (54) 124 Xu, J., see Ohkuma, S. (54) 133

Yamada, N., see Hirao, T. (54) 49 Yamamoto, H., see Hamada, S. (54) 199 Yamamoto, T., see Hamada, S. (54) 199 Yamamoto, Y., see Hirao, T. (54) 49 Yang, W., see Loh, H.H. (54) 321 Yokoya, S., see Kanazawa, K.-y. (54) 316 Yoshikawa, S., see Hamada, S. (54) 199 Yoshimura, T., see Kamata, T. (54) 219 Young, A.B., see Friberg, I.K. (54) 298 Yuhki, N., see Kamata, T. (54) 219

Ziółkowska, B., Przewłocka, B., Mika, J., Łabuz, D. and Przewłocki, R. Evidence for Fos involvement in the regulation of proenkephalin and prodynorphin gene expression in the rat hippocampus (54) 243